ALBANIAN JOURNAL OF MATHEMATICS Volume 5, Number 3, Pages 151–164 ISSN: 1930-1235; (2011)

GENERALIZED HYERS-ULAM STABILITY OF DERIVATIONS IN PROPER LIE CQ*-ALGEBRAS

G.Z. ESKANDANI Faculty of Mathematical Science, University of Tabriz, Tabriz, Iran Email: zamani@tabrizu.ac.ir

J. M. RASSIAS Pedagogical Department E.E., National and Capodistrian University of Athens, Email: jrassias@primedu.uoa.gr

> R. ZARGHAMI Faculty of Mathematical Science, University of Tabriz, Tabriz, Iran, Email: zarghami@tabrizu.ac.ir

ABSTRACT. In this paper, we obtain the general solution and the generalized Hyers-Ulam stability for the following functional equation

$$f(\frac{\sum_{i=1}^{m} x_i}{m}) + \sum_{\substack{i=1\\i \neq j}}^{m} f(\frac{x_j - x_i}{m}) = f(x_j).$$

This is applied to investigate derivations and their stability in proper Lie $CQ^\ast\text{-}$ algebras.

1. INTRODUCTION AND PRELIMINARIES

Ulam [42] gave a talk before the Mathematics Club of the University of Wisconsin in which he discussed a number of unsolved problems. Among these problems was the following question concerning the stability of homomorphisms.

©2011 Aulona Press (Albanian J. Math.)

²⁰¹⁰ Mathematics Subject Classification. 17B40, 39B52, 47N50, 47L60, 46B03.

 $Key\ words\ and\ phrases.$ Generalized Hyers-Ulam stability, proper Lie $CQ^*\text{-algebra}$, Lie derivation.

Let $(G_1, *)$ be a group and let (G_2, \diamond, d) be a metric group with the metric $d(\cdot, \cdot)$. Given $\epsilon > 0$, does there exist a $\delta(\epsilon) > 0$ such that if a mapping $f : G_1 \to G_2$ satisfies the inequality

$$d(f(x * y), f(x) \diamond f(y)) < \delta$$

for all $x, y \in G_1$, then there is a homomorphism $T: G_1 \to G_2$ with

$$d(f(x), T(x)) < \epsilon$$

for all $x \in G_1$?

If the answer is affirmative, we say that the equation of homomorphism T(xy) = T(x)T(y) is stable. The concept of stability for a functional equation arises when we replace the functional equation by an inequality which acts as a perturbation of the equation. Thus the stability question of functional equations is that how do the solutions of the inequality differ from those of the given functional equation?

Hyers [18] considered the case of approximately additive mappings $f: E \to E'$, where E and E' are Banach spaces and f satisfies *Hyers inequality*

$$\|f(x+y) - f(x) - f(y)\| \le \epsilon$$

for all $x, y \in E$. It was shown that the limit

$$L(x) = \lim_{n \to \infty} \frac{f(2^n x)}{2^n}$$

exists for all $x \in E$ and that $L: E \to E'$ is the unique additive mapping satisfying

$$\|f(x) - L(x)\| \le \epsilon.$$

Hyers' theorem was generalized by Aoki [3] for additive mappings and independently by Th.M. Rassias [36] for linear mappings by considering an *unbounded Cauchy difference*. In 1994, a generalization of Th.M. Rassias' theorem was obtained by Găvruta [15]. J.M. Rassias [31]-[34] generalized Hyers result. During the last two decades, a number of papers and research monographs have been published on various generalizations and applications of the generalized Hyers-Ulam stability to a number of functional equations and mappings (see [11]-[13], [20], [24]-[28],[30], [37]-[39]). We also refer the readers to the books [1], [10], [19], [21] and [37].

We recall some basic facts concerning quasi *-algebras.

Definition 1.1. Let A be a linear space and A_0 be a *-algebra contained in A as a subspace. We say that A is a *quasi* *-*algebra* over A_0 if

- (i) the right and left multiplications of an element of A and an element of A_0 are always defined and linear;
- (*ii*) $x_1(x_2a) = (x_1x_2)a$, $(ax_1)x_2 = a(x_1x_2)$ and $x_1(ax_2) = (x_1a)x_2$ for all $x_1, x_2 \in A_0$ and all $a \in A$;

(*iii*) an involution *, which extends the involution of A_0 , is defined in A with the property $(ab)^* = b^*a^*$, whenever the multiplication is defined.

Quasi *-algebras [22, 23] arise in natural way as completions of locally convex *-algebras whose multiplication is not jointly continuous; in this case one has to deal with topological quasi *-algebras.

A quasi *-algebra (A, A_0) is called *topological* if a locally convex topology τ on A is given such that:

- (i) the involution $a \mapsto a^*$ is continuous for each $a \in A$,
- (*ii*) the mappings $a \mapsto ab$ and $a \mapsto ba$ are continuous for each $a \in A$ and $b \in A_0$,
- (*iii*) A_0 is dense in $A[\tau]$.

Throughout this paper, we suppose that a locally convex quasi *-algebra (A, A_0) is complete. For an overview on partial *-algebra and related topics we refer to [2].

In a series of papers [4], [5], [6], [7] many authors have considered a special class of quasi *-algebras, called proper CQ^* -algebras, which arise as completions of C^* -algebras. They can be introduced in the following way:

Definition 1.2. Let A be a Banach module over the C^* -algebra A_0 with involution * and C^* -norm $\|.\|_0$ such that $A_0 \subset A$. We say that (A, A_0) is a proper CQ^* -algebra if

- (i) A_0 is dense in A with respect to its norm $\|.\|$;
- (*ii*) $(ab)^* = b^*a^*$ whenever the multiplication is defined;
- $(iii) ||y||_0 = \max\{\sup_{a \in A, ||a|| \le 1} ||ay||, \sup_{a \in A, ||a|| \le 1} ||ya|| \} \text{ for all } y \in A_0.$

A proper CQ^* -algebra (A, A_0) is said to have a unit e if there exists an element $e \in A_0$ such that ae = ea = a for all $a \in A$. In this paper we will always assume that the proper CQ^* -algebra under consideration have an identity.

Definition 1.3. A proper CQ^* -algebra (A, A_0) , endowed with a bilinear multiplication $[,]: (A \times A_0) \cup (A_0 \times A) \to A$, called the bracket, which satisfies two simple properties:

- (i) $[x_1, x_2] = -[x_2, x_1]$ for all $(x_1, x_2) \in (A \times A_0) \cup (A_0 \times A);$
- (*ii*) $[x_1, [x_2, x_3]] = [[x_1, x_2], x_3] + [x_1, [x_2, x_3]]$ for all $x_1, x_2, x_3 \in A_0$

is called a proper Lie CQ^* -algebra.

Definition 1.4. Let (A, A_0) be a proper Lie CQ^* -algebras. A \mathbb{C} -linear mapping $\delta : A_0 \to A$ is called a *Lie derivation* if

$$\delta([z, x]) = [\delta(z), x] + [z, \delta(x)]$$

for all $x, z \in A_0$ (see [28]).

Throughout this paper, we assume that m and j are fixed positive integers with $m \ge 2$.

In this paper, we obtain the general solution and the generalized Ulam-Hyers stability for the following functional equation

(1.1)
$$f(\frac{\sum_{i=1}^{m} x_i}{m}) + \sum_{\substack{i=1\\i\neq j}}^{m} f(\frac{x_j - x_i}{m}) = f(x_j)$$

where m is a fixed positive integer with $m \ge 2$. This is applied to investigate derivations and their stability on proper Lie CQ^* -algebras.

2. Solution of functional equation (1.1)

Throughout this section, let both X and Y be real vector spaces. We here present the general solution of (1.1).

Theorem 2.1. A mapping $f : X \to Y$ satisfies (1.1) if and only if the mapping $f : X \to Y$ is additive.

We first assume that the mapping $f: X \to Y$ satisfies (1.1). Setting $x_j = x$ and $x_i = 0$ for all $1 \le i \le m$ and $i \ne j$ in (1.1), we get

(2.1)
$$f\left(\frac{x}{m}\right) = \frac{1}{m}f(x)$$

for all $x \in X$. Setting $x_j = x$, $x_{j+1} = y$ and $x_i = 0$ for $i \neq j, j+1$ in (1.1) and using (2.1), we get

(2.2)
$$f\left(\frac{x+y}{m}\right) + f\left(\frac{x-y}{m}\right) = \frac{2}{m}f(x)$$

for all $x, y \in X$. Replacing x and y by mx and my in (2.2), we get

(2.3)
$$f(x+y) + f(x-y) = 2f(x)$$

for all $x, y \in X$. Setting y = x in (2.3), we get

$$(2.4) f(2x) = 2f(x)$$

for all $x \in X$. Replacing x by $\frac{x+y}{2}$ and y by $\frac{x-y}{2}$ in (2.3), and using (2.4) we get

$$f(x+y) = f(x) + f(y)$$

for all $x, y \in X$. So the mapping $f : X \to Y$ is additive.

Conversely, let the mapping $f: X \to Y$ be additive. By a simple computation, one can show that the mapping f satisfies the functional equation (1.1).

3. Stability of derivation on proper Lie CQ^* -algebras

Throughout this section, assume that (A, A_0) is a proper Lie CQ^* -algebra with C^* -norm $\|.\|_{A_0}$ and norm $\|.\|_A$. For convenience, we use the following abbreviation for a given mapping $f: \underbrace{A_0 \times A_0 \times \ldots \times A_0}_{\longrightarrow} \to A$

$$D_{\mu}f(x_{1},...,x_{m}) := f\left(\frac{\sum_{i=1}^{m} \mu x_{i}}{m}\right) + \sum_{\substack{i=1\\i\neq j}}^{m} f\left(\frac{\mu x_{j} - \mu x_{i}}{m}\right) - \mu f(x_{j})$$

for all $x_1, \dots, x_m \in A_0$, where $\mu \in \mathbb{T}^1 := \{\mu \in \mathbb{C} : |\mu| = 1\}$. We will use the following lemma:

Lemma 3.1. [29] Let $f : A_0 \to A$ be an additive mapping such that $f(\mu x) = \mu f(x)$ for all $x \in A_0$ and all $\mu \in \mathbb{T}^1$. Then the mapping f is \mathbb{C} -linear.

Theorem 3.2. Let $\varphi : \underbrace{A_0 \times A_0 \times \ldots \times A_0}_{m-times} \to [0,\infty) \text{ and } \psi : A_0 \times A_0 \to [0,\infty) \text{ be}$

mappings such that

(3.1)
$$\lim_{n \to \infty} \frac{1}{m^n} \varphi(m^n x_1, ..., m^n x_m) = 0$$

(3.2)
$$\lim_{n \to \infty} \frac{1}{m^{2n}} \psi(m^n x_1, m^n x_2) = 0,$$

(3.3)
$$\widetilde{\varphi_j}(x) := \sum_{i=1}^{\infty} \frac{1}{m^i} \varphi(0, ..., \underbrace{m^i x}_{j \ th}, ...0) < \infty$$

for all $x, x_1, \dots, x_m \in A_0$. Suppose that $f : A_0 \to A$ is a mapping such that

(3.4)
$$||D_{\mu}f(x_1,...,x_m)||_A \le \varphi(x_1,...,x_m),$$

(3.5)
$$\left\| f([x_1, x_2]) - [f(x_1), x_2] - [x_1, f(x_2)] \right\|_A \le \psi(x_1, x_2)$$

for all $x_1, \dots, x_m \in A_0$ and all $\mu \in \mathbb{T}^1$. Then there exists a unique Lie derivation $\delta : A_0 \to A$ such that

(3.6)
$$||f(x) - \delta(x)||_A \le \widetilde{\varphi_j}(x)$$

for all $x \in A_0$.

Letting $\mu = 1$, $x_j = mx$ and $x_i = 0$ for all $1 \le i \le m$ with $i \ne j$ in (3.4), we get (3.7) $\|f(mx) - mf(x)\|_A \le \varphi(0, ..., \underbrace{mx}_{j \ th}, ...0)$

for all $x \in A_0$. Replacing x by $m^n x$ in (3.7) and dividing both sides of (3.7) by m^{n+1} , we get

(3.8)
$$\left\|\frac{1}{m^{n+1}}f(m^{n+1}x) - \frac{1}{m^n}f(m^nx)\right\|_A \le \frac{1}{m^{n+1}}\varphi(0,...,\underbrace{m^{n+1}x}_{j\,th},...,0)$$

for all $x \in A_0$ and all non-negative integers n. Hence

(3.9)
$$\begin{aligned} \left\|\frac{1}{m^{n+1}}f(m^{n+1}x) - \frac{1}{m^k}f(m^kx)\right\|_A &\leq \sum_{i=k}^n \left\|\frac{1}{m^{i+1}}f(m^{i+1}x) - \frac{1}{m^i}f(m^ix)\right\|_A \\ &\leq \sum_{i=k+1}^{n+1}\frac{1}{m^i}\varphi(0,...,\underbrace{m^ix}_{j\,th},...,0) \end{aligned}$$

for all $x \in A_0$ and all non-negative integers n and k with $n \ge k$. Therefore, we conclude from (3.3) and (3.9) that the sequence $\{\frac{1}{m^n}f(m^nx)\}_n$ is a Cauchy sequence in A for all $x \in A_0$. Since A is complete, the sequence $\{\frac{1}{m^n}f(m^nx)\}_n$ converges in A for all $x \in A_0$. So one can define the mapping $\delta : A_0 \to A$ by

(3.10)
$$\delta(x) := \lim_{n \to \infty} \frac{1}{m^n} f(m^n x)$$

for all $x \in A_0$. Letting k = 0 and passing the limit $n \to \infty$ in (3.9), we get (3.6). Now, we show that δ is a \mathbb{C} -linear mapping. It follows from (3.1), (3.4) and (3.10) that

$$\|D_1\delta(x_1,...,x_m)\|_A = \lim_{n \to \infty} \frac{1}{m^n} \|D_1f(m^n x_1,...,m^n x_m)\|_A$$
$$\leq \lim_{n \to \infty} \frac{1}{m^n} \varphi(m^n x_1,...,m^n x_m) = 0$$

for all $x_1, \dots, x_m \in A_0$. So the mapping δ satisfies (1.1). By Theorem 2.1, the mapping δ is additive.

Letting $x_j = mx$ and $x_i = 0$ for all $1 \le i \le m$ with $i \ne j$ in (3.4), we get

(3.11)
$$\|mf(\mu x) - \mu f(mx)\|_A \le \varphi(0, ..., \underbrace{mx}_{j \ th}, ..., 0)$$

for all $x \in A_0$. Replacing x by $m^n x$ in (3.11) and dividing both sides of (3.11) by m^{n+1} , we get

(3.12)
$$\begin{aligned} \left\| \frac{1}{m^n} f(\mu m^n x) - \frac{\mu}{m^{n+1}} f(m^{n+1} x) \right\|_A \\ &\leq \frac{1}{m^{n+1}} \varphi(0, ..., \underbrace{m^{n+1} x}_{j \ th}, ..., 0) \end{aligned}$$

for all $x \in A_0$ and all non-negative integers n. Passing the limit $n \to \infty$ in (3.12) and using (3.1) and (3.10), we get

$$\delta(\mu x) = \mu \delta(x)$$

for all $\mu \in \mathbb{T}^1$ and for all $x \in A_0$. So by Lemma 3.1, we infer that the mapping $\delta : A_0 \to A$ is \mathbb{C} -linear. To prove the uniqueness of δ , let $\delta' : A_0 \to A$ be another additive mapping satisfying (3.6). It follows from (3.6) and (3.10) that

$$\begin{split} \|\delta(x) - \delta'(x)\|_A &= \lim_{n \to \infty} \frac{1}{m^n} \|f(m^n x) - \delta'(m^n x)\|_A \\ &\leq \lim_{n \to \infty} \frac{1}{m^n} \widetilde{\varphi_j}(m^n x) = 0 \end{split}$$

for all $x \in A_0$. So $\delta = \delta'$.

It follows from (3.2), (3.5) and (3.10) that

$$\begin{split} \left\| \delta([x_1, x_2]) - [\delta(x_1), x_2] - [x_1, \delta(x_2)] \right\|_A \\ &= \lim_{n \to \infty} \frac{1}{m^{2n}} \left\| f(m^{2n}[x_1, x_2]) - [f(m^n x_1), m^n x_2] - [m^n x_1, f(m^n x_2)] \right\|_A \\ &\leq \lim_{n \to \infty} \frac{1}{m^{2n}} \psi(m^n x_1, m^n x_2) = 0 \end{split}$$

for all $x_1, x_2 \in A_0$. So

$$\delta([x_1, x_2]) = [\delta(x_1), x_2] + [x_1, \delta(x_2)]$$

for all $x_1, x_2 \in A_0$. Hence the mapping $\delta : A_0 \to A$ is a unique Lie derivation satisfying (3.6).

Corollary 3.3. Let $\delta, \alpha_1, \alpha_2, s_1, s_2, \{\theta_i\}_{i=1}^m$ and $\{r_i\}_{i=1}^m$ be non-negative real numbers such that $0 < s_1, s_2 < 2$, and $0 < r_i < 1$ for all $1 \le i \le m$. Suppose that $f: A_0 \to A$ is a mapping such that

$$\begin{aligned} \|D_{\mu}f(x_{1},...,x_{m})\|_{A} &\leq \delta + \sum_{i=1}^{m} \theta_{i} \|x_{i}\|_{A_{0}}^{r_{i}}, \\ \|f([x_{1},x_{2}]) - [f(x_{1}),x_{2}] - [x_{1},f(x_{2})]\|_{A} &\leq \delta + \alpha_{1} \|x_{1}\|_{A_{0}}^{s_{1}} + \alpha_{2} \|x_{2}\|_{A_{0}}^{s_{2}}, \end{aligned}$$

for all $x_1, \dots, x_m \in A_0$ and all $\mu \in \mathbb{T}^1$. Then there exists a unique Lie derivation $\delta : A_0 \to A$ such that

$$||f(x) - \delta(x)||_A \le \frac{\delta}{m-1} + \gamma(x)$$

for all $x \in A_0$, where

$$\gamma(x) := \min_{1 \le i \le m} \left\{ \frac{\theta_i m^{r_i}}{m - m^{r_i}} \|x\|_{A_0}^{r_i} \right\}.$$

Corollary 3.4. Let $\delta, \alpha_1, \alpha_2, \alpha_3, s_1, s_2$ and $\{r_i\}_{i=1}^m$ be non-negative real numbers such that $s_1 + s_2 < 2$ and $0 < \sum_{i=1}^m r_i < 1$ for all $1 \le i \le m$. Suppose that $f: A_0 \to A$ is a mapping such that

$$\|D_{\mu}f(x_1,...,x_m)\|_A \le \delta + \sum_{i=1}^m \|x_i\|_{A_0}^{r_i} + \prod_{i=1}^m \|x_i\|_{A_0}^{r_i},$$

 $\|f([x_1, x_2]) - [f(x_1), x_2] - [x_1, f(x_2)]\|_A \le \delta + \alpha_1 \|x_1\|_{A_0}^{s_1} + \alpha_2 \|x_2\|_{A_0}^{s_2} + \alpha_3 \|x_1\|_{A_0}^{s_1} \|x_2\|_{A_0}^{s_2},$

for all $x_1, \dots, x_m \in A_0$ and all $\mu \in \mathbb{T}^1$. Then there exists a unique Lie derivation $\delta : A_0 \to A$ such that

$$\|f(x) - \delta(x)\|_A \le \frac{\delta}{m-1} + \tau(x)$$

for all $x \in A_0$, where

$$\tau(x) := \min_{1 \le i \le m} \left\{ \frac{m^{r_i}}{m - m^{r_i}} \|x\|_{A_0}^{r_i} \right\}.$$

Note that the mixed "product-sum" function was introduced by J. M. Rassias in 2008-09 ([8, 9, 16, 17, 40, 41]).

Theorem 3.5. Let $\Phi: \underbrace{A_0 \times A_0 \times \ldots \times A_0}_{m-times} \to [0,\infty)$ and $\Psi: A_0 \times A_0 \to [0,\infty)$ be

mappings such that

(3.13)

$$\lim_{n \to \infty} m^n \Phi\left(\frac{x_1}{m^n}, ..., \frac{x_m}{m^n}\right) = 0,$$

$$\lim_{n \to \infty} m^{2n} \Psi\left(\frac{x_1}{m^n}, \frac{x_2}{m^n}\right) = 0,$$

$$\widetilde{\Phi_j}(x) := \sum_{i=0}^{\infty} m^i \Phi(0, ..., \underbrace{\frac{x_i}{m^i}}_{j \ th}, ..., 0) < \infty$$

for all $x, x_1, \dots, x_m \in A_0$. Suppose that $f : A_0 \to A$ is a mapping such that

$$\|D_{\mu}f(x_1,...,x_m)\|_A \le \Phi(x_1,...,x_m),$$

$$\|f([x_1,x_2]) - [f(x_1),x_2] - [x_1,f(x_2)]\|_A \le \Psi(x_1,x_2)$$

for all $x_1, \dots, x_m \in A_0$ and all $\mu \in \mathbb{T}^1$. Then there exists a unique Lie derivation $\delta: A_0 \to A$ such that

(3.14)
$$||f(x) - \delta(x)||_A \le \widetilde{\Phi_j}(x)$$

for all $x \in A_0$.

Similarly to the proof of Theorem 3.2, we have

(3.15)
$$\|f(mx) - mf(x)\|_A \le \Phi(0, ..., \underbrace{mx}_{j \ th}, ...0)$$

for all $x \in A_0$. Replacing x by $\frac{x}{m^{n+1}}$ in (3.15) and multiplying both sides of (3.15) to m^n , we get

$$\left\| m^{n+1} f\left(\frac{x}{m^{n+1}}\right) - m^n f\left(\frac{x}{m^n}\right) \right\|_A \le m^n \Phi\left(0, \dots, \underbrace{\frac{x}{m^n}}_{j \ th}, \dots 0\right)$$

for all $x \in A_0$ and all non-negative integers n. Hence

$$(3.16) \qquad \left\| m^{n+1} f\left(\frac{x}{m^{n+1}}\right) - m^k f\left(\frac{x}{m^k}\right) \right\|_A \le \sum_{i=k}^n \left\| m^{i+1} f\left(\frac{x}{m^{i+1}}\right) - m^i f\left(\frac{x}{m^i}\right) \right\|_A$$
$$\le \sum_{i=k}^n m^i \Phi\left(0, \dots, \underbrace{\frac{x}{m^i}}_{j \ th}, \dots 0\right)$$

for all $x \in A_0$ and all non-negative integers n and k with $n \ge k$. Therefore the sequence $\{m^n f(\frac{x}{m^n})\}$ is a Cauchy sequence in A for all $x \in A_0$. Since A is complete,

the sequence $\{m^n f(\frac{x}{m^n})\}$ converges in A for all $x \in A_0$. So one can define the mapping $\delta : A_0 \to A$ by

$$\delta(x) := \lim_{n \to \infty} m^n f\left(\frac{x}{m^n}\right)$$

for all $x \in A_0$. Letting k = 0 and passing the limit $n \to \infty$ in (3.16), we get (3.14).

The rest of the proof is similar to the proof of Theorem 3.2.

Corollary 3.6. Let $\alpha_1, \alpha_2, s_1, s_2, \{\theta_i\}_{i=1}^m$ and $\{r_i\}_{i=1}^m$ be non-negative real numbers such that $s_1, s_2 > 2$ and $r_i > 1$ for all $1 \le i \le m$. Suppose that $f : A_0 \to A$ is a mapping such that

$$\begin{split} \|D_{\mu}f(x_{1},...,x_{m})\|_{A} &\leq \sum_{i=1}^{m} \theta_{i} \|x_{i}\|_{A_{0}}^{r_{i}}, \\ \|f([x_{1},x_{2}]) - [f(x_{1}),x_{2}] - [x_{1},f(x_{2})]\|_{A} &\leq \alpha_{1} \|x_{1}\|_{A_{0}}^{s_{1}} + \alpha_{2} \|x_{2}\|_{A_{0}}^{s_{2}}, \end{split}$$

for all $x_1, \dots, x_m \in A_0$ and all $\mu \in \mathbb{T}^1$. Then there exists a unique Lie derivation $\delta : A_0 \to A$ such that

$$||f(x) - \delta(x)||_A \le \Gamma(x)$$

for all $x \in A_0$, where

$$\Gamma(x) := \min_{1 \le i \le m} \big\{ \frac{\theta_i m^{r_i}}{m^{r_i} - 1} \|x\|_{A_0}^{r_i} \big\}.$$

Corollary 3.7. Let $\alpha_1, \alpha_2, \alpha_3, s_1, s_2$ and $\{r_i\}_{i=1}^m$ be non-negative real numbers such that $s_1, s_2 > 2$ and $r_i > 1$ for all $1 \le i \le m$. Suppose that $f : A_0 \to A$ is a mapping such that

$$\|D_{\mu}f(x_1,...,x_m)\|_A \le \sum_{i=1}^m \|x_i\|_{A_0}^{r_i} + \prod_{i=1}^m \|x_i\|_{A_0}^{r_i},$$

 $\|f([x_1, x_2]) - [f(x_1), x_2] - [x_1, f(x_2)]\|_A \le \alpha_1 \|x_1\|_{A_0}^{s_1} + \alpha_2 \|x_2\|_{A_0}^{s_2} + \alpha_3 \|x_1\|_{A_0}^{s_1} \|x_2\|_{A_0}^{s_2},$

for all $x_1, \dots, x_m \in A_0$ and all $\mu \in \mathbb{T}^1$. Then there exists a unique Lie derivation $\delta : A_0 \to A$ such that

$$||f(x) - \delta(x)||_A \le \Delta(x)$$

for all $x \in A_0$, where

$$\Delta(x) := \min_{1 \le i \le m} \left\{ \frac{m^{r_i}}{m^{r_i} - m} \|x\|_{A_0}^{r_i} \right\}.$$

4. Subadditive mapping and stability of Eq. (1.1)

Next, using some idea of [35], we are going to establish other theorems about the stability of Eq. (1.1)

We call that a subadditive mapping is a mapping $\varphi : A \to B$, having a domain A and a codomain (B, \leq) that are both closed under addition, with the following property:

$$\varphi(x+y) \le \varphi(x) + \varphi(y)$$

for all $x, y \in X$. Now we say that a mapping $\varphi : X \to Y$ is contractively subadditive if there exists a constant L with 0 < L < 1 such that

$$\varphi(x+y) \le L[\varphi(x) + \varphi(y)]$$

for all $x, y \in X$. Therefore φ satisfies the following properties $\varphi(mx) \leq mL\varphi(x)$ and so $\varphi(m^n x) \leq (mL)^n \varphi(x)$, for all $x \in X$ and all positive integer $m \geq 2$. Similarly, we say that a mapping $\varphi : A \to B$ is expansively superadditive if there exists a constant L with 0 < L < 1 such that

$$\varphi(x+y) \geq \frac{1}{L}[\varphi(x)+\varphi(y)]$$

for all $x, y \in X$. Therefor φ satisfies the following properties $\varphi(x) \leq \frac{L}{m}\varphi(mx)$ and so $\varphi(\frac{x}{m^n}) \leq (\frac{L}{m})^n \varphi(x)$, for all $x \in X$ and all positive integer $m \geq 2$.

Theorem 4.1. Let $\varphi : \underbrace{A_0 \times A_0 \times \ldots \times A_0}_{m-times} \to [0,\infty)$ be a contractively subadditive with the constant L and $\psi : A_0 \times A_0 \to [0,\infty)$ be a mapping such that

(4.1)
$$\lim_{n \to \infty} \frac{1}{m^{2n}} \psi(m^n x_1, m^n x_2) = 0,$$

for all $x_1, x_2 \in A_0$. Suppose that $f : A_0 \to A$ is a mapping such that

(4.2)
$$||D_{\mu}f(x_1,...,x_m)||_A \le \varphi(x_1,...,x_m),$$

(4.3)
$$\left\| f([x_1, x_2]) - [f(x_1), x_2] - [x_1, f_1(x_2)] \right\|_A \le \psi(x_1, x_2)$$

for all $x_1, \dots, x_m \in A_0$ and all $\mu \in \mathbb{T}^1$. Then there exists a unique Lie derivation $\delta : A_0 \to A$ such that

(4.4)
$$||f(x) - \delta(x)||_A \le \frac{L}{1 - L}\varphi(0, ..., \underbrace{x}_{j \ th}, ...0)$$

for all $x \in X$.

Letting $\mu = 1$, $x_j = mx$ and $x_i = 0$ for all $1 \le i \le m$ with $i \ne j$ in (4.2), we get

(4.5)
$$||f(mx) - mf(x)||_A \le \varphi(0, ..., \underbrace{mx}_{jth}, ...0)$$

for all $x \in A_0$.

Replacing x by $m^n x$ in (4.5) and dividing both sides of (4.5) by m^{n+1} , we get

(4.6)
$$\left\|\frac{1}{m^{n+1}}f(m^{n+1}x) - \frac{1}{m^n}f(m^nx)\right\|_A \le \frac{1}{m^{n+1}}\varphi(0,...,\underbrace{m^{n+1}x}_{j\,th},...,0) \le \frac{(mL)^{n+1}}{m^{n+1}}\varphi(0,...,\underbrace{x}_{j\,th},...,0) \le L^{n+1}\varphi(0,...,\underbrace{x}_{j\,th},...,0)$$

for all $x \in A_0$ and all non-negative integers n. Hence

(4.7)
$$\begin{aligned} \left\|\frac{1}{m^{n+1}}f(m^{n+1}x) - \frac{1}{m^k}f(m^kx)\right\|_A &\leq \sum_{i=k}^n \left\|\frac{1}{m^{i+1}}f(m^{i+1}x) - \frac{1}{m^i}f(m^ix)\right\|_A \\ &\leq \sum_{i=k+1}^{n+1} L^i\varphi(0,...,\underbrace{x}_{ith},...,0) \end{aligned}$$

for all $x \in A_0$ and all non-negative integers n and k with $n \ge k$. Therefore, we conclude from and (4.7) that the sequence $\{\frac{1}{m^n}f(m^nx)\}$ is a Cauchy sequence in A for all $x \in A_0$. Since A is complete, the sequence $\{\frac{1}{m^n}f(m^nx)\}$ converges in A for all $x \in A_0$. So one can define the mapping $\delta : A_0 \to A$ by

(4.8)
$$\delta(x) := \lim_{n \to \infty} \frac{1}{m^n} f(m^n x)$$

for all $x \in A_0$. Letting k = 0 and passing the limit $n \to \infty$ in (4.7), we get (4.4). Now, we show that δ is a \mathbb{C} -linear mapping. It follows from (4.8) that

$$\|D_1\delta(x_1,...,x_m)\|_A = \lim_{n \to \infty} \frac{1}{m^n} \|D_1f(m^n x_1,...,m^n x_m)\|_A$$

$$\leq \lim_{n \to \infty} \frac{1}{m^n} \varphi(m^n x_1,...,m^n x_m)$$

$$\leq \lim_{n \to \infty} L^n \varphi(x_1,...,x_m) = 0$$

for all $x_1, \dots, x_m \in A_0$. So the mapping δ satisfies (1.1). By Lemma 2.1, the mapping δ is additive.

Letting $x_j = mx$ and $x_i = 0$ for all $1 \le i \le m$ with $i \ne j$ in (4.2), we get

(4.9)
$$||mf(\mu x) - \mu f(mx)||_A \le \varphi(0, ..., \underbrace{mx}_{j \ th}, ..., 0)$$

for all $x \in A_0$. Replacing x by $m^n x$ in (4.9) and dividing both sides of (4.9) by m^{n+1} , we get

(4.10)
$$\begin{aligned} \left\| \frac{1}{m^n} f(\mu m^n x) - \frac{\mu}{m^{n+1}} f(m^{n+1} x) \right\|_A \\ &\leq \frac{1}{m^{n+1}} \varphi(0, ..., \underbrace{m^{n+1} x}_{j \ th}, ..., 0) \end{aligned}$$

for all $x \in A_0$ and all non-negative integers *n*. Passing the limit $n \to \infty$ in (4.10) and using (4.8), we get

$$\delta(\mu x) = \mu \delta(x)$$

for all $\mu \in \mathbb{T}^1$ and for all $x \in A_0$. So by Lemma 3.1, we infer that the mapping $\delta : A_0 \to A$ is \mathbb{C} -linear. To prove the uniqueness of δ , let $\delta' : A_0 \to A$ be another

additive mapping satisfying (4.4). It follows from (4.8) that

$$\begin{aligned} \|\delta(x) - \delta'(x)\|_A &= \lim_{n \to \infty} \frac{1}{m^n} \|f(m^n x) - \delta'(m^n x)\|_A \\ &\leq \lim_{n \to \infty} \frac{L^{n+1}}{1 - L} \varphi(0, \dots, \underbrace{x}_{jth}, \dots 0) = 0 \end{aligned}$$

for all $x \in A_0$. So $\delta = \delta'$.

The rest of the proof is similar to the proof of Theorem 3.2.

Corollary 4.2. Let θ be non-negative real number and $f : A_0 \to A$ be a mapping for which

$$\|D_{\mu}f(x_1, ..., x_m)\|_A \le \theta$$
$$\left\|f([x_1, x_2]) - [f(x_1), x_2] - [x_1, f_1(x_2)]\right\|_A \le \theta$$

 θ

for all $x_1, ..., x_m \in A_0$. Then there exists a unique Lie derivation $\delta : A_0 \to A$ such that

(4.11)
$$||f(x) - \delta(x)||_A \le \theta$$

for all $x \in A_0$.

The proof follows from Theorem 4.1 by taking

$$\varphi(x_1, \dots, x_m) := \theta$$

for all $x_1, \ldots, x_m \in A_0$.

Replacing contractively subadditive by expansively superadditive in Theorem 4.1, one can obtain the following theorem:

Theorem 4.3. Let $\varphi : \underbrace{A_0 \times A_0 \times \ldots \times A_0}_{m-times} \to [0,\infty)$ be a expansively superadditive with the constant L and $\psi : A_0 \times A_0 \to [0,\infty)$ be a mapping such that

(4.12)
$$\lim_{n \to \infty} m^{2n} \psi\left(\frac{x_1}{m^n}, \frac{x_2}{m^n}\right) = 0,$$

for all $x_1, x_2 \in A_0$. Suppose that $f : A_0 \to A$ is a mapping such that

(4.13)
$$||D_{\mu}f(x_1,...,x_m)||_A \le \varphi(x_1,...,x_m),$$

(4.14)
$$\left\| f([x_1, x_2]) - [f(x_1), x_2] - [x_1, f_1(x_2)] \right\|_A \le \psi(x_1, x_2)$$

for all $x_1, \dots, x_m \in A_0$ and all $\mu \in \mathbb{T}^1$. Then there exists a unique Lie derivation $\delta : A_0 \to A$ such that

(4.15)
$$||f(x) - \delta(x)||_A \le \frac{1}{1 - L}\varphi(0, ..., \underbrace{x}_{jth}, ...0)$$

for all $x \in X$.

STABILITY OF DERIVATIONS

References

- J. Aczél, J. Dhombres, Functional Equations in Several Variables, Cambridge University Press, 1989.
- [2] J.P. Antoine, A. Inoue and C. Trapani, Partial *-Algebras and Their Operator Realizations, Kluwer, Dordrecht, 2002.
- [3] T. Aoki, On the stability of the linear transformation in Banach spaces, J. Math. Soc. Japan 2 (1950) 64-66.
- [4] F. Bagarello, A. Inoue and C. Trapani, Some classes of topological quasi *-algebras, Proc. Amer. Math. Soc. 129 (2001), 2973–2980.
- [5] F. Bagarello and C.Trapani, States and representations of CQ*-algebras, Ann. Inst. H. Poincaré 61 (1994), 103–133.
- [6] F. Bagarello and C. Trapani, CQ*-algebras: Structure properties, Publ. Res. Inst. Math. Sci. 32 (1996), 85–116.
- [7] F. Bagarello and C. Trapani, Morphisms of certain Banach C^{*}-modules, Publ. Res. Inst. Math. Sci. 36 (2000), 681–705.
- [8] H. X. Cao, J. R. Lv and J. M. Rassias, Superstability for generalized module left derivations and generalized module derivations on a banach module (I), Journal of Inequalities and Applications, Volume 2009, Art. ID 718020, 1–10.
- [9] H. X. Cao, J. R. Lv and J. M. Rassias, Superstability for generalized module left derivations and generalized module derivations on a banach module(II), J. Pure. Appl. Math. 10 (2009), Issue 2, 1–8.
- [10] P. Czerwik, Functional Equations and Inequalities in Several Variables, World Scientific Publishing Company, New Jersey, Hong Kong, Singapore and London, 2002.
- [11] G. Z. Eskandani, On the Hyers-Ulam-Rassias stability of an additive functional equation in quasi-Banach spaces, J. Math. Anal. Appl., 345 (2008), 405-409.
- [12] G. Z. Eskandani, P. Gavruta, J. M. Rassias, and R. Zarghami, Generalized Hyers-Ulam stability for a general mixed functional equation in quasi-β-normed spaces, Mediterr. J. Math. 8 (2011), 331-348.
- [13] G. Z. Eskandani, H. Vaezi and Y. N. Dehghan, Stability of a mixed additive and quadratic functional equation in non-Archimedean Banach modules, 11 (2010), 1309–1324.
- [14] Z. Gajda, On stability of additive mappings, Intern. J. Math. Sci. 14 (1991), 431–434.
- [15] P. Găvruta, A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings, J. Math. Anal. Appl. 184 (1994), 431–436.
- [16] M.E. Gordji and H. Khodaei, On the generalized Hyers-Ulam-Rassias stability of quadratic functional equations, Abstract and Applied Analysis, Volume 2009, Art. ID 923476, 1–11, Doi:10.1155/2009/923476.
- [17] M.E. Gordji, S.Zolfaghari, J.M. Rassias and M.B. Savadkouhi, Solution and Stability of a Mixed type Cubic and Quartic functional equation in Quasi-Banach spaces, Abstract and Applied Analysis, Volume 2009, Art. ID 417473, 1–14, Doi:10.1155/2009/417473.
- [18] D.H. Hyers, On the stability of the linear functional equation, Proc. Nat. Acad. Sci. USA 27 (1941), 222–224.
- [19] D.H. Hyers, G. Isac and Th.M. Rassias, Stability of Functional Equations in Several Variables, Birkhäuser, Basel, 1998.
- [20] K. Jun and Y. Lee, On the Hyers-Ulam-Rassias stability of a Pexiderized quadratic inequality, Math. Inequal. Appl. 4 (2001), 93–118.
- [21] S. M. Jung, Hyers-Ulam-Rassias Stability of Functional Equations in Mathimatical Analysis, Hadronic Press, Palm Harbor, 2001.
- [22] G. Lassner, Topological algebras and their applications in quantum statistics, Wiss.Z. KMU, Leipzig, Math.-Nat.R. 6, 572–595 (1981); Algebras of Unbounded Operators and Quantum Dynamics, Physica, 124A, (1984) 471–480.
- [23] G. Lassner and G.A. Lassner, Quasi *-algebras and Twisted Product, Publ. RIMS, Kyoto Univ. 25, (1989) 279–299.
- [24] F. Moradlou, H. Vaezi and G.Z. Eskandani, Hyers-Ulam-Rassias stability of a quadratic and additive functional equation in quasi-Banach spaces. Mediterr. J. Math. 6 (2009), no. 2, 233-248.
- [25] M.S. Moslehian, Ternary derivations, stability and physical aspects, Acta Appl. Math. 100, (2008) 187–199.

- [26] A. Najati and G.Z. Eskandani, Stability of derivations on proper Lie CQ*-algebras. Commun. Korean Math. Soc. 24 (2009), 5-16.
- [27] A. Najati and G.Z. Eskandani, Stability of a mixed additive and cubic functional equation in quasi-Banach spaces, J. Math. Anal. Appl. 342 (2008) 1318-1331.
- [28] C. Park, Homomorphisms between Lie JC*-algebras and Cauchy-Rassias stability of Lie JC*-algebra derivations, J. Lie Theory 15 (2005), 393–414.
- [29] C. Park, Homomorphisms between Poisson JC*-algebras, Bull. Braz. Math. Soc. 36 (2005), 79–97.
- [30] C. Park and Th.M. Rassias, Homomorphisms and derivations in proper JCQ*-triples, J. Math. Anal. Appl. 337 (2008), 1404–1414.
- [31] J.M. Rassias, On approximation of approximately linear mappings by linear mappings, Journal of Functional Analysis, 46(1982), 126-130.
- [32] J. M. Rassias, On approximation of approximately linear mappings by linear mappings, Bulletin des Sciences Mathematiques, 108(1984), 445-446.
- [33] J. M. Rassias, Solution of a problem of Ulam, Journal of Approximation Theory, 57(1989), 268-273,.
- [34] J.M. Rassias, Solution of a stability problem of Ulam, Discussiones Mathematicae, 12(1992), 95-103.
- [35] J.M. Rassias and H.M. Kim, Generalized Hyers-Ulam stability for general additive functional equations in quasi-β-normed spaces, J. Math. Anal. Appl. 356 (2009), 302–309.
- [36] Th.M. Rassias, On the stability of the linear mapping in Banach spaces, Proc. Amer. Math. Soc. 72 (1978), 297–300.
- [37] Th.M. Rassias (ed.), Functional Equations and Inequalities, Kluwer Academic Publishers, Dordrecht, Boston, London, 2000.
- [38] Th.M. Rassias, On a modified Hyers-Ulam sequence, J. Math. Anal. Appl. 158 (1991) 106– 113.
- [39] Th.M. Rassias, Problem 16; 2, Report of the 27th International Symp. on Functional Equations, Aequationes Math. 39 (1990) 292–293.
- [40] K. Ravi, M. Arunkumar and J. M. Rassias, Ulam stability for the orthogonally general Euler-Lagrange type functional equation, Intern. J. Math. Stat. 3 (A08)(2008), 36–46.
- [41] K. Ravi, J. M. Rassias, M. Arunkumar, and R. Kodandan, Stability of a generalized mixed type additive, quadratic, cubic and quartic functional equation, J. Pure. Appl. Math. 10(2009), Issue 4, Article 114, 1–29.
- [42] S.M. Ulam, A Collection of the Mathematical Problems, Interscience Publ., New York, 1960.